

# Advanced Quantum Field Theory

## Exam problems

1. Find the free massive propagator  $\Delta(x - x')$  for a  $(1 + 1)$ -dimensional spacetime and study the behavior for  $x - x'$  timelike (inside the lightcone) and  $x - x'$  spacelike (outside the lightcone).
2. For a real scalar field with interaction  $\lambda\varphi^4/4!$ , draw all the connected Feynman diagram contributions to the two-point function  $G^{(2)}$  and the four-point function  $G^{(4)}$  up to order  $\lambda^3$ .
3. Consider a four dimensional real scalar field with the Lagrangian

$$\mathcal{L} = -\frac{1}{2}\partial^\mu\varphi\partial_\mu\varphi - \frac{1}{2}m^2\varphi^2 - \frac{1}{4!}\lambda\varphi^4.$$

Compute the lowest-order corrections to the propagator and compute the  $Z$ -factors of the quadratic terms in the Lagrangian in the  $\overline{MS}$  scheme.

4. Yukawa theory is defined as a 4D theory of a Dirac fermion and real scalar field defined by the Lagrangian

$$\begin{aligned}\mathcal{L} = & i\bar{\psi}\not{\partial}\psi - m\bar{\psi}\psi - \frac{1}{2}\partial^\mu\varphi\partial_\mu\varphi - \frac{1}{2}M^2\varphi^2 \\ & + ig\varphi\bar{\psi}\gamma_5\psi - \frac{1}{4!}\lambda\varphi^4.\end{aligned}$$

Derive the fermion-loop correction to the scalar propagator. Show that there is an extra minus sign for a fermion loop as compared to a scalar loop.

5. Assume that starting with the Maxwell Lagrangian  $\mathcal{L}_{\text{Max}}$  in 4D we fix the gauge with the gauge fixing condition

$$\partial^\mu A_\mu + \sigma A^\mu A_\mu = 0,$$

where  $\sigma$  is an arbitrary real constant. How would the gauge fixing term and the ghost Lagrangian look like? Show that there are now self-interactions among the photons. What is the gauge propagator? What is the ghost propagator? What are the ghost-gauge vertices? Calculate the divergent one-loop contributions to the two-point function of the gauge field. Feynman gauge  $\xi = 1$  may be assumed.

6. In pure Yang-Mills theory fix the gauge using the axial gauge condition

$$n^\mu A_\mu^a = 0$$

for  $n^\mu$  a fixed four-vector. Find the gluon and ghost propagators and the ghost-gluon interaction vertices.

7. In four dimension, calculate the contribution to the Yang-Mills beta function from a non-self-interacting complex scalar field transforming in the representation  $R$  of the gauge group, coupled to a non-abelian external gauge field

$$\mathcal{L} = -D^\mu \phi^\dagger D_\mu \phi - m^2 \phi^\dagger \phi.$$

8. Consider performing the path integral for a scalar field theory in the presence of a background field  $\bar{\varphi}(x)$ . We define

$$e^{i\widetilde{W}[J, \bar{\varphi}]} = \int \mathcal{D}\varphi e^{iS[\varphi + \bar{\varphi}] + i \int d^d x J \varphi}.$$

Clearly  $\widetilde{W}[J, 0]$  is the original  $W[J]$ . We also define the quantum action in the presence of the background field

$$\widetilde{\Gamma}[\varphi, \bar{\varphi}] = \widetilde{W}[J, \bar{\varphi}] - \int d^d x J \varphi,$$

where now  $J(x)$  is the solution of

$$\frac{\delta}{\delta J(x)} \widetilde{W}[J, \bar{\varphi}] = \varphi(x).$$

Show that  $\widetilde{\Gamma}[\varphi, 0]$  is equal to the original quantum action  $\Gamma[\varphi]$  and that

$$\Gamma[\varphi + \bar{\varphi}] = \widetilde{\Gamma}[\varphi + \bar{\varphi}, 0] = \widetilde{\Gamma}[\varphi, \bar{\varphi}],$$

which means that we can calculate the original quantum action by calculating vacuum graphs in the background action

$$\Gamma[\bar{\varphi}] = \widetilde{\Gamma}[\bar{\varphi}, 0] = \widetilde{\Gamma}[0, \bar{\varphi}].$$

Confirm this by computing the one-loop contribution to the two-point function for a real scalar field in 4D with a  $\frac{1}{4!}\varphi^4$  potential using this method and in the standard way and compare the results.