

User manual for the SCPH -program

by
Petros Souvatzis

Department of Physics



Uppsala University 2008

Contents

1	Introduction	3
2	Theoretical Background	4
2.1	The harmonic lattice and its limitations	4
2.2	The SCAILD method	6
2.3	Theoretical appendix	8
3	Setting up a SCAILD calculation	9
3.1	Overview of the SCAILD calculation	10
3.2	The input files	11
3.2.1	The INPHON-file	11
3.2.2	The POSCAR-file	13
3.2.3	The INPUTFLJ-file	13
3.2.4	The DIPOLEINTERACT-file	16
3.2.5	The KRAFTER-file	17
3.2.6	The KRAFTER.1, , KRAFTER.n-files	17
3.2.7	The QPOINTS and the SYMOP -files	17
3.2.8	The MEMORY-file	18
3.3	The output files	18
3.3.1	The MEMORY-file	18
3.3.2	The POSCARTEMP1-file	19
3.3.3	The DISPLACEMENTS-file	19
3.3.4	The FREQ-file	19
3.3.5	The DOS-file	19
3.3.6	The DOSPROJ-file	19
3.3.7	The CONVERGENCE-file	20
3.3.8	The QPOINTS and SYMOP-file	20
3.3.9	The BLANDAD-file	21
4	Model-potential calculations	22
4.1	The MSV-potential	22
4.2	The ZRP-potential	23
4.3	The EAM-potential	23
4.4	The PAS-potential	24
4.5	The PIN-potential	25
4.6	The PAA-potential	26

Chapter 1

Introduction

In this manual/user guide describes how to operate the scph-program in which the self consistent ab initio lattice dynamical method has been implemented (SCAILD) [1]. The program is written in a way so that it can be easily be interfaced with the VASP package [2], and with the phon-program of Dario Alfe [3]. Those of you who have used VASP and phon will thus find it a "peace of cake" to learn how to use the scph package.

There are also three different directories accompanying this manual with all the input files needed to do three different SCAILD calculations. These directories are `examples/bcc.Ti/`, `examples/hcp.Ti/` and `examples/CsI/` and it might be a good idea to take a look at these files when reading this manual.

Chapter 2

Theoretical Background

Self consistent phonon calculations are a natural extension of the theory of the harmonic lattice, and we initiate our methodological description by first presenting the most important features of this theory. The Hamiltonian of lattice dynamical system can be expressed as

$$\mathcal{H} = \mathcal{H}_h + \mathcal{V}_{anh} \quad (2.1)$$

where

$$\mathcal{H}_h = \sum_{\mathbf{R}, \sigma} \frac{\mathbf{P}_{\mathbf{R}\sigma}^2}{2M_\sigma} + \frac{1}{2} \sum_{\mathbf{R}, \sigma} \sum_{\mathbf{R}', \sigma'} \mathbf{U}_{\mathbf{R}\sigma} \bar{\bar{\Phi}}^{\sigma\sigma'}(\mathbf{R} + \mathbf{b}_\sigma - \mathbf{R}' - \mathbf{b}_{\sigma'}) \mathbf{U}_{\mathbf{R}'\sigma'}, \quad (2.2)$$

$$\mathcal{V}_{anh} = \frac{1}{3!} \sum_{\mathbf{R}, \sigma} \sum_{\mathbf{R}', \sigma'} \sum_{\mathbf{R}'', \sigma''} \sum_{\alpha\beta\gamma} \Phi_{\alpha\beta\gamma}^{\sigma\sigma'\sigma''}(\mathbf{R} + \mathbf{b}_\sigma, \mathbf{R}' + \mathbf{b}_{\sigma'}, \mathbf{R}'' + \mathbf{b}_{\sigma''}) \mathbf{U}_{\mathbf{R}\sigma\alpha} \mathbf{U}_{\mathbf{R}'\sigma'\beta} \mathbf{U}_{\mathbf{R}''\sigma''\gamma} + \dots, \quad (2.3)$$

are the harmonic and anharmonic parts of the Hamiltonian, respectively. Here \mathbf{R} are the equilibrium lattice positions of the atoms, \mathbf{b}_σ the atomic positions relative to the primitive Bravais lattice, $\mathbf{U}_{\mathbf{R}\sigma}$ the displacements of the atoms, $\mathbf{P}_{\mathbf{R}\sigma}$ the momentum of the atoms, M_σ the atomic mass, $\bar{\bar{\Phi}}^{\sigma\sigma'}$ the inter-atomic force constant matrices and $\Phi_{\alpha,\beta,\gamma}^{\sigma\sigma'\sigma''}$ the tensor describing the third order anharmonic contribution to the potential energy. In order to make the notation more transparent, in the following sections the notation of a monoatomic lattice will be adapted without any loss of generality.

2.1 The harmonic lattice and its limitations

Diagonalizing the dynamical matrix

$$\bar{\bar{D}}(\mathbf{k}) = \frac{1}{M} \sum_{\mathbf{R}} \bar{\bar{\Phi}}(\mathbf{R}) e^{-i\mathbf{k}\mathbf{R}}, \quad (2.4)$$

for each wave vector \mathbf{k} in the first Brillouin zone one finds the eigenvalues $\omega_{\mathbf{k}s}$ and eigenvectors $\epsilon_{\mathbf{k}s}$ of different phonon modes (longitudinal or transverse) labeled by the symbol s ,

N being the number of atoms. Introducing the canonical phonon coordinates $Q_{\mathbf{k}s}$ and $P_{\mathbf{k}s}$, the displacements $\mathbf{U}_{\mathbf{R}}$ and the kinetic energy operators $\mathbf{P}_{\mathbf{R}}$ can be expressed as

$$\mathbf{U}_{\mathbf{R}} = \frac{1}{\sqrt{MN}} \sum_{\mathbf{k},s} Q_{\mathbf{k}s} \epsilon_{\mathbf{k}s} e^{i\mathbf{k}\mathbf{R}} \quad (2.5)$$

$$\mathbf{P}_{\mathbf{R}} = \frac{1}{\sqrt{MN}} \sum_{\mathbf{k},s} P_{\mathbf{k}s} \epsilon_{\mathbf{k}s} e^{i\mathbf{k}\mathbf{R}} \quad (2.6)$$

allowing a separation of the harmonic Hamiltonian \mathcal{H}_h of the crystal into the Hamiltonians of $3N$ independent harmonic oscillators.

$$\mathcal{H}_h = \sum_{\mathbf{k}s} \frac{1}{2} (P_{\mathbf{k}s}^2 + \omega_{\mathbf{k}s}^2 Q_{\mathbf{k}s}^2). \quad (2.7)$$

The thermodynamic average of the operators $Q_{\mathbf{k}s}^\dagger Q_{\mathbf{k}s}$ determines the mean-square atomic displacements and is given by

$$\langle Q_{\mathbf{k}s}^\dagger Q_{\mathbf{k}s} \rangle = \frac{\hbar}{\omega_{\mathbf{k}s}} \left[\frac{1}{2} + n \left(\frac{\hbar \omega_{\mathbf{k}s}}{k_B T} \right) \right], \quad (2.8)$$

where $n(x) = 1/(e^x - 1)$ is the Planck function. In the classical limit, i.e for sufficiently high temperatures, the operators $(1/\sqrt{M})Q_{\mathbf{k}s}$ are replaced by real numbers,

$$\mathcal{R}_{\mathbf{k}s} \equiv \pm \sqrt{\frac{\langle Q_{\mathbf{k}s}^\dagger Q_{\mathbf{k}s} \rangle}{M}}. \quad (2.9)$$

Calculating the gradient of the potential energy in Eqn. (2.2) with respect to the atomic displacements gives the restoring force

$$\mathbf{F}_{\mathbf{R}} = - \sum_{\mathbf{R}'} \bar{\Phi}(\mathbf{R} - \mathbf{R}') \mathbf{U}_{\mathbf{R}'}. \quad (2.10)$$

Fourier transforming Eqn.(2.10) and substituting $\mathbf{U}_{\mathbf{R}}$ with the expression in Eqn. (2.5) gives

$$\mathbf{F}_{\mathbf{k}} = - \sum_s M \omega_{\mathbf{k}s}^2 \mathcal{R}_{\mathbf{k}s} \epsilon_{\mathbf{k}s}. \quad (2.11)$$

Finally, using the orthogonality of the eigenvectors $\epsilon_{\mathbf{k}s}$ the phonon frequencies can be expressed as

$$\omega_{\mathbf{k}s} = \left[- \frac{1}{M} \frac{\epsilon_{\mathbf{k}s} \mathbf{F}_{\mathbf{k}}}{\mathcal{R}_{\mathbf{k}s}} \right]^{1/2}. \quad (2.12)$$

The equations discussed so far can be solved for dynamically stable materials, where the excitation of any individual phonon mode, i.e a finite $Q_{\mathbf{k}s}$, will lead to an increase of the total energy E_{tot} of the system, i.e $dE_{tot}/dQ_{\mathbf{k}s} \geq 0$. For dynamically unstable materials however, there exists one or several phonon modes (\mathbf{k}, s) such that any finite $Q_{\mathbf{k}s}$ for these modes will result in a decrease in the total energy of the lattice. In this situation the equations discussed so far can not be used straight forwardly since they result in imaginary phonon frequencies. This represents a situation where the lattice under consideration spontaneously

shifts atomic planes and/or atomic positions so that a new crystal structure lowers the total energy.

The only type of anharmonicity that can be taken into account by the above formalism is the type of anharmonicity responsible for the thermal expansion. This is a weaker kind of anharmonicity that most people are aware of is present in all solids. Qualitatively it can be described by the simple argument that longer bonds are weaker, so the frequencies are lower, so the entropy is higher. To capture the anharmonicity associated to the thermal expansion quasiharmonic DFT works very well. Here the change of phonon frequencies is almost entirely due to thermal expansion. However, whenever strong anharmonicity appears quasiharmonic phonons may show unstable modes which cannot be stabilized by the thermal expansion. Thus the interaction between phonons must be included in order to correctly describe the lattice dynamics of the systems. In the following section it will be explained how the interaction between phonons can be taken into account by means of the SCAILD method.

2.2 The SCAILD method

To understand how the SCAILD incorporates the effect of anharmonicity in the calculations of phonon frequencies it is instructive to substitute the lattice displacements in the Hamiltonian (2.1) with the decomposition of $U_{\mathbf{R}}$ given by Eqn. (2.5), to obtain the following expression for the Hamiltonian

$$\mathcal{H} = \frac{1}{2} \sum_{\mathbf{k}s} \left[\mathcal{P}_{\mathbf{k}s}^2 + \omega_{\mathbf{k}s}^2 \left(1 + \frac{1}{3} \sum_{\mathbf{k}_1, \mathbf{k}_2} \sum_{s_1, s_2} \mathcal{A}(\mathbf{k}, \mathbf{k}_1, \mathbf{k}_2, s, s_1, s_2) \frac{\mathcal{Q}_{\mathbf{k}_1 s_1} \mathcal{Q}_{\mathbf{k}_2 s_2}}{\mathcal{Q}_{\mathbf{k}s} \omega_{\mathbf{k}s}^2} + \dots \right) \mathcal{Q}_{\mathbf{k}s}^2 \right] \quad (2.13)$$

where

$$\mathcal{A}(\mathbf{k}, \mathbf{k}_1, \mathbf{k}_2, s, s_1, s_2) = \frac{1}{(MN)^{3/2}} \sum_{\mathbf{R}, \mathbf{R}_1, \mathbf{R}_2} \sum_{\alpha, \beta, \gamma} \Phi_{\alpha\beta\gamma}(\mathbf{R}, \mathbf{R}_1, \mathbf{R}_2) \epsilon_{\mathbf{k}s\alpha} \epsilon_{\mathbf{k}_1 s_1 \beta} \epsilon_{\mathbf{k}_2 s_2 \gamma} e^{i(\mathbf{R}\mathbf{k} + \mathbf{R}_1 \mathbf{k}_1 + \mathbf{R}_2 \mathbf{k}_2)} \quad (2.14)$$

The full Hamiltonian of Eqn. (2.13) is contrary to the Harmonic problem not separable into N independent Hamiltonians. Nevertheless by replacing the operators $\mathcal{Q}_{\mathbf{k}s}$ appearing in the curved brackets of Eqn. (2.13) with $\sqrt{M}\mathcal{R}_{\mathbf{k}s}$ the following mean-field Hamiltonian can be constructed

$$\mathcal{H}^{MF} = \sum_{\mathbf{k}s} \frac{1}{2} (\mathcal{P}_{\mathbf{k}s}^2 + \bar{\omega}_{\mathbf{k}s}^2 \mathcal{Q}_{\mathbf{k}s}^2) \quad (2.15)$$

where

$$\bar{\omega}_{\mathbf{k}s}^2 = \omega_{\mathbf{k}s}^2 \left(1 + \frac{\sqrt{M}}{2} \sum_{\mathbf{k}_1, \mathbf{k}_2} \sum_{s_1, s_2} \mathcal{A}(\mathbf{k}, \mathbf{k}_1, \mathbf{k}_2, s, s_1, s_2) \frac{\mathcal{R}_{\mathbf{k}_1 s_1} \mathcal{R}_{\mathbf{k}_2 s_2}}{\mathcal{R}_{\mathbf{k}s} \omega_{\mathbf{k}s}^2} + \dots \right) \quad (2.16)$$

The set of equations 2.8-2.9 and 2.15-2.16 can, once the interaction terms $\mathcal{A}(\mathbf{k}, \mathbf{k}_1, \mathbf{k}_2, s, s_1, s_2)$, and an initial guess in terms of normal modes have been provided, be solved self consistently.

It follows from experience of calculations of anharmonic effects in metals [10, 11] that usually odd-order terms are dominant for close packed stiff lattices, such as Ir metal [10]. At the same time, soft-mode phonon behavior in essentially anharmonic bcc metals, such as

Ca and Sr, is determined by the even-order terms [11]. The phonon damping is connected mainly with three-phonon processes and cannot be taken into account in the framework of self-consistent phonon picture. In this sense, our approach is reminiscent the quasiparticle self-consistent GW approximation [12] where only real part of the electron self-energy is taken into account.

In the SCAILD scheme Equations (2.8)-(2.9) and (2.15)-(2.16) are solved by first calculating a starting guess for the phonon dispersions by means of a standard supercell calculation, see e.g Ref. [13]. The phonon frequencies corresponding to \mathbf{k} -vectors commensurate with the supercell are then used to calculate the atomic displacements through Eqn. (2.5) and Eqns. (2.8)-(2.9). Here it should be noted that the signs of the amplitudes $\mathcal{A}_{\mathbf{k}s}$ (see Eqn.(2.9)), should be chosen randomly, with equal probabilities for + and -. It should also be stressed that in the first iterations of the calculation unstable modes may be present, i.e there may be modes with $\omega_{\mathbf{k}s}^2 < 0$. In order to calculate the Fourier amplitude $\mathcal{R}_{\mathbf{k}s}$ for these modes by means Eqns. (2.8)-(2.9) the frequencies have to be guessed. In these situations the absolute value $|\omega_{\mathbf{k}s}|$ of the imaginary frequencies have been used as guesses of the real frequencies.

The forces induced by the displacements $U_{\mathbf{R}}$ can be calculated by any standard ab initio method. From the Fourier transform of the atomic forces a new set of frequencies $\bar{\omega}_{\mathbf{k}s}^2$ are calculated through Eqn. (2.12) (in the appendix we describe how one from Eqn. (2.12) can extract $\bar{\omega}_{\mathbf{k}s}^2$) The symmetries of the different \mathbf{k} -vectors are restored by

$$\Omega_{\mathbf{k}s}^2 = \frac{1}{m_{\mathbf{k}}} \sum_{S \in \mathcal{S}(\mathbf{k})} \bar{\omega}_{S^{-1}\mathbf{k}S}^2, \quad (2.17)$$

where $\mathcal{S}(\mathbf{k})$ is the symmetry group of the wave vector \mathbf{k} , and m_k the number of elements of the group. The mean value of all iterations supplies a new set of frequencies,

$$\omega_{\mathbf{k}s}^2(N_I) = \frac{1}{N_I} \sum_{i=1}^{N_I} \Omega_{\mathbf{k}s}^2(i), \quad (2.18)$$

where $\Omega_{\mathbf{k}s}(i)$, $i = 1, \dots, N_I$ are the symmetry restored frequencies from all iterations. Here N_I is the number of iterations. The new set of frequencies calculated in (2.18) determine a new set of displacements used to calculate a new set of forces. In Fig. 2.1 a schematic outline is shown of the different steps performed in a SCAILD calculation. Philosophically our approach is similar to Born's self consistent phonon theory, with the main difference being that we consider a direct force calculation from a super cell with Hellman-Feynman forces calculated from density functional theory.




Figure 2.1: (color online). Schematic outline over the different tasks performed in a self-consistent phonon calculation.

2.3 Theoretical appendix

This appendix shows how the eigenvalues $\bar{\omega}_{\mathbf{k}s}^2$ of the mean field Hamiltonian (2.15) can be retained from Eqn. (2.12). Using the sum convention for the coordinate indexes the forces can be expressed as

$$F_{\mathbf{R}j} = - \sum_{\mathbf{R}'} \bar{\Phi}_{j\alpha}(\mathbf{R} - \mathbf{R}') U_{\mathbf{R}'\alpha} - \frac{1}{2} \sum_{\mathbf{R}', \mathbf{R}''} \Phi_{j\alpha\beta}(\mathbf{R}, \mathbf{R}', \mathbf{R}'') U_{\mathbf{R}'\alpha} U_{\mathbf{R}''\beta} + \dots \quad (2.19)$$

By replacing the operators $\mathcal{Q}_{\mathbf{k}s}/\sqrt{M}$ with the real numbers $\mathcal{A}_{\mathbf{k}s}$ in the normal mode expansion of $\mathbf{U}_{\mathbf{R}}$, the Fourier transform of the forces can be expressed as

$$F_{\mathbf{k}j} = - \sum_s M \omega_{\mathbf{k}s}^2 \epsilon_{\mathbf{k}s j} \mathcal{R}_{\mathbf{k}s} - \frac{1}{2N^{3/2}} \sum_{\mathbf{k}_1, \mathbf{k}_2} \sum_{s_1, s_2} \Phi_{j\alpha\beta}(\mathbf{k}, \mathbf{k}_1, \mathbf{k}_2) \epsilon_{\mathbf{k}_1 s_1 \alpha} \epsilon_{\mathbf{k}_2 s_2 \beta} \mathcal{R}_{\mathbf{k}_1 s_1} \mathcal{R}_{\mathbf{k}_2 s_2} + \dots \quad (2.20)$$

Finally multiplying (2.20) with $-\epsilon_{\mathbf{k}s j}/(\mathcal{R}_{\mathbf{k}s} M)$ and summing over j gives

$$-\frac{\mathbf{F}_{\mathbf{k}} \epsilon_{\mathbf{k}s}}{M \mathcal{R}_{\mathbf{k}s}} = \omega_{\mathbf{k}s}^2 \left(1 + \frac{\sqrt{M}}{2} \sum_{\mathbf{k}_1, \mathbf{k}_2} \sum_{s_1, s_2} \mathcal{A}(\mathbf{k}, \mathbf{k}_1, \mathbf{k}_2, s, s_1, s_2) \frac{\mathcal{R}_{\mathbf{k}_1 s_1} \mathcal{R}_{\mathbf{k}_2 s_2}}{\mathcal{R}_{\mathbf{k}s} \omega_{\mathbf{k}s}^2} + \dots \right). \quad (2.21)$$

where $\mathcal{A}(\mathbf{k}, \mathbf{k}_1, \mathbf{k}_2, s, s_1, s_2)$ is given by (2.14). Now from 2.21 becomes obvious that the left hand side of (2.12) is equivalent to $\bar{\omega}_{\mathbf{k},s}$.

Chapter 3

Setting up a SCAILD calculation

In order to set up a SCAILD calculation one needs, apart from the scph-program, a first principles program that can calculate inter-atomic forces. Any first principles program can of course be used for this purpose, however since the input- and output-format of the scph-program follows that of VASP, it will be assumed in this manual that the ab initio program is VASP. The scph-program and the ab initio program are interfaced by means of a simple shell script. The structure of the script is the same as the structure found in the schematic picture presented in Fig. 2.1. Translating this picture into a simple bash script one would obtain the following: (Observe that the small numbers to the right of each active line of the script are not part of the script, they are just used for reference)

```
#!/bin/sh
#First displacement
cp POSCAR.START1 POSCAR 1
./vasp 2
./force_extract 3
cp FORCESI KRAFTER.1 4

...

#n:th displacement
cp POSCAR.STARTn POSCAR 5
./vasp 6
./force_extract 7
cp FORCESI KRAFTER.n 8

#Here we run scph and generate starting guess displacements:
cp POSCAR.REF POSCAR 9
./scph 10
```

```

for i in 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 11
do 12
cp POSCARTEMP1 POSCAR13
./vasp 14
./force_extract 15
cp FORCESI KRAFTER 16
cp POSCAR_REF POSCAR17
./scph18
done19

```

3.1 Overview of the SCAILD calculation

In this section a short overview of the different tasks performed in a SCAILD calculation will be discussed. All along this discussion references will be made to the generic bash script that was presented in the very beginning of this chapter.

The lines 1-4 and 5-8 in the script is the part of the SCAILD calculation where the initial phonon dispersion is calculated. This initial calculation is what is referred to as the starting guess calculation (see STEP 1 in Fig. 2.1). Here the files `POSCAR.START1` and `POSCAR.STARTn` contain the atomic positions of the supercell. In these files all but one of the atoms are placed in their equilibrium positions. The number of files/initial displacements needed to calculate the initial phonon dispersion depend upon the symmetry of the crystal. In order to generate the files `POSCAR.START1` - `POSCAR.STARTn`, the `phon`-program [3] has been used. At lines 1-2 and 5-6 the forces induced by the displaced atoms in `POSCAR.START1` - `POSCAR.STARTn` are calculated. At lines 3 and 7 the perl script `force_extract` is used to extract the forces in the `OUTCAR`-files into the file `FORCESI`. The `FORCESI`-files are then copied to the files `KRAFTER.1` - `KRAFTER.n` on lines 4 and 8 so that they later on may be recognized and used by the `scph` program on line 10.

On line 9 the file `POSCAR_REF` containing the equilibrium coordinates of the supercell is copied to the file `POSCAR`, which is to be used as input by the `scph` program on line 10. At line 10 the `scph` program uses the files `KRAFTER.1` - `KRAFTER.n` and `POSCAR` (The following files are also used: `INPHON`, `INPUTFLJ`, `QPOINTS` and `SYMOP`, but they will be discussed in the next section) to calculate the force constant matrices $\bar{\Phi}(\mathbf{R})$, the initial phonon dispersion, the atomic displacements $\mathbf{U}_{\mathbf{R}}$ through Eqn. 2.5, 2.8 and 2.9 and the new atomic positions $\mathbf{R} + \mathbf{U}_{\mathbf{R}}$ (see STEP 2 in Fig. 2.1). The new atomic positions are then written to the file `POSCARTEMP1` by the `scph` program.

At lines 11-19 the script enters the main loop of the SCAILD calculation, where self consistency is to be obtained (see STEP 2, 3, 4 and 5 in Fig. 2.1). At lines 13-15 the forces $\mathbf{F}_{\mathbf{R}}$ acting on the atoms positioned at $\mathbf{R} + \mathbf{U}_{\mathbf{R}}$ are calculated (see STEP 3 in Fig. 2.1). At lines 16-18 the forces stored in the file `KRAFTER` are used by the `scph` program to calculate a new set of phonon frequencies through Eqn. 2.12, 2.17 and 2.18 (see STEP 4 and 5 in Fig. 2.1). New atomic positions $\mathbf{R} + \mathbf{U}_{\mathbf{R}}$ are then calculated by the `scph` program and written to the file `POSCARTEMP1`. The file `POSCARTEMP1` is then used at lines 13 and 14 to calculate a new set of forces. (i.e back to step 2 in Fig. 2.1).

3.2 The input files

The input files of the `scph`-program are the following:

```
INPHON
POSCAR
INPUTFLJ
DIPOLEINTERACT
KRAFTER
KRAFTER.1, . . . , KRAFTER.n
QPOINTS
SYMOP
MEMORYFILE
```

3.2.1 The INPHON-file

This is the same `INPHON`-file as the one used by the `phon`-program [3]. Here only those features important for the `scph`-program will be commented, for a more complete description of the file please look into the manual of the `phon`-program [3]. Below an example of a typical `INPHON`-file is given:

```
# symmetryze force constant matrix
# LSymm=.FALSE.

# number of ions types and masses
NTYPES = 1; MASS = 44.956 1

# generate superlattice
LSUPER = .F.; NDIM = 4 4 4; DISP = 1002

# free energy calculation
LFREE = .F.; TEMPERATURE = 16733

# q points section
LRECIP = .T.4
ND = 4; NPOINTS = 1005
QI = 0.0 0.0 0.0 \6
0.5 -0.5 0.5 \7
0.25 0.25 0.25 \8
0.0 0.0 0.0
QF = 0.5 -0.5 0.5 \9
0.25 0.25 0.25 \10
0.0 0.0 0.0 \11
0.0 0.0 0.512

# density of states
LGAMMA = .TRUE.13
QA = -100; QB = 100 ; QC = 10014
DOSIN = 0; DOSEND = 10; DOSSTEP = 0.05; DOSSMEAR = .05115
# write force constant matrix
```

LFORCEOUT = .T.¹⁶

In the following the contents of the above example INPHON-file will be referenced to by the small number appearing to the right of each active line.

Line (1):

Here the number of different atomic species are entered together with the different masses. This specific file refers to a SCAILD calculation of bcc Sc. For example if this file were to be used in a calculation of ZrO₂ the corresponding line would read:

NTYPES = 2; MASS = 91.224 15.9994

Line (2):

Here the flag **LSUPER** should always be set to false when the INPHON-file is used by the **scph**-program. However if the INPHON-file is used by the **phon**-program [3], then setting this flag to true will make the **phon**-program create a supercell by increasing the primitive cell given by the **POSCAR**-file along the three primitive lattice vectors. The increase/multiplication of the cell is given by the array **NDIM**. The resulting supercell will be written to the file **SPOSCAR**. Whenever the INPHON-file is to be used by the **scph**-program the array stored in **NDIM** should correspond to the supercell size used. In the above example **NDIM = 4 4 4** corresponds to a SCAILD calculation with a 64-atom supercell. The parameter **DISP** is unimportant and can be set to an arbitrary value when the INPHON-file is to be used by the **scph**-program.

Line (3):

Whenever the INPHON-file is to be used by the **scph**-program the flag **LFREE** should be set to false. The parameter **TEMPERATURE** is more or less self explanatory and it is the temperature T appearing in Eqn. 2.8 and in the calculation of the phonon contribution to the free energy given by

$$F^{phon}(T) = \int_0^\infty d\omega g(\omega) \left[\frac{\hbar\omega}{2} + k_B T \ln(1 - e^{-\hbar\omega/k_B T}) \right], \quad (3.1)$$

where $g(\omega)$ is the phonon density of states.

Line (4-12):

These lines specify between which points in reciprocal space the phonon dispersion $\omega_{\mathbf{k},s}$ should be calculated. The output is written to the file **FREQ**. For a more detailed description of this input please have a look in the manual of the **phon**-program [3].

Line (13):

Please have a look in the manual of the **phon**-program [3].

Line (14-15):

These lines determines how the phonon density of states $g(\omega)$ is to be calculated. The parameters **QA**, **QB** and **QC** is the mesh in reciprocal space used to calculate the density of states $g(\omega)$. If the INPHON-file is used by the **phon**-program the minus sign in **QA** "tells" the **phon**-program that the q-point mesh already exists and is present in the file **QPOINTS**. If the first mesh number, i.e **QA**, is positive the **phon**-program will calculate the q-points used to calculate the density of states $g(\omega)$ and write them to the file **QPOINTS**.

If the INPHON-file is to be used by the **scph**-program then the parameters **QA**, **QB** and **QC** have

the same meaning as described above. However in this case the QA parameter should always be negative. In the event that the QPOINTS-file doesn't exist the scph-program will at the first iteration (STEP 1 in the SCAILD calculation, see Fig. 2.1) generate the QPOINTS-file. The parameters DOSIN and DOSEND specifies the frequency range in THz in which the density of states should be calculated. The parameter DOSSTEP is the maximum resolution of each state in THz. The parameter DOSSMEAR is the gaussian smearing in THz used to smear each state.

Line (16):

If this flag is true the force constant matrix $\bar{\bar{\Phi}}$ will be written to the file HARMONIC.

3.2.2 The POSCAR-file

This file contain the structural information of the system. It has exactly the same format as the format used by the VASP package [2]. The POSCAR-file used by the scph-program contains the equilibrium atomic positions of the atoms of the supercell. The dimensions of the supercell stored in the POSCAR-file should be the same as specified by the array NDIM in the INPHON-file. The simplest way to obtain a desired POSCAR-file is to just take the SPOSCAR-file generated with the phon-program [3].

3.2.3 The INPUTFLJ-file

This file "tells" the scph-program whether or not the inter-atomic forces are to be calculated externally by some ab initio code, or internally by means of model potentials. Furthermore the INPUTFLJ-file also specifies the number of initial displacements, the amplitude and direction of these displacements and which atoms that have been displaced in the initial calculation (STEP 1 in Fig. 2.1). Below an example of an INPUTFLJ-file is given:

```
POT ABI1
NDISP 12
DISPL 1 0.0 0.01 0.01 1 0.0 0.0 0.02 1 0.0 0.0 0.02 1 0.0 0.0 0.023.....
RM 3.23234
BETA -1.13635
X1 1.300006
X2 1.650007
B1 0.013312008
B2 -0.245730009
B3 1.904700010
B4 -8.05410011
C6 20.2160012
C8 -30.806013
C10 28.50600000014
EPS 11.28516908015
RCUT 30.016
RK 1.00000000 1.05000000 1.55000000 1.60000000 1.65000000 1.70000000 1.750000017
AK -38.4084159 36.9240324 -6.06706900 8.45632500 -4.68932080 7.95571790 -5.644938918
AZ -0.44417136 1.07926027 -0.67936231 -14.90219
SAMPLING NGAUSS20
```

```

NGAUSSAMPLE 121
MAXITTER 6022
DSITTER 1023
SUPERSAFE .FALSE.24
MAXAMP 1.025

```

Line(1):

The parameter POT "tells" the `scph`-program whether or not the inter-atomic forces are to be calculated externally by some ab initio code, or internally by means of model potentials. If the parameter is equal to `ABI` the inter-atomic forces will be calculated externally, and the lines 4-19 will not be used since they only refer to model-potential parameter data. If the parameter POT equals any of the following three letter combinations `MSV`, `ZRP`, `EAM`, `PAS` or `PIN` the inter-atomic forces will be calculated internally from model-potentials. In chapter 4 the model potentials will be discussed in greater detail.

Line(2):

The parameter `NDISP` equals the number of initial single atom displacements needed to obtain the starting guess phonon frequencies (STEP1, in Fig. 2.1). This version of the `scph` can only manage a maximum of 6 initial displacements.

Line(3):

The array `DISPL` contains 6 blocks with 4 elements each. The first element in each block refers to the atom in `POSCAR`-file that has been displaced. The following three elements in each block describe the direction and amplitude of the displacement in direct coordinates. In the above example of an `INPUTFLJ`-file one initial displacement is used (`NDISP=1`), the atom being displaced is the atom with the topmost coordinate appearing in the `POSCAR`-file, and the displacement is described by the vector 0.0000 0.01 0.01 i.e

`DISPL 1 0.0 0.01 0.01, . . .`. In the above example the 20 following elements of the array `DISPL`, i.e

```
1 0.0 0.0 0.02  1 0.0 0.0 0.02  1 0.0 0.0 0.02 ....
```

are not used since `NDISP` equals 1. Nevertheless the number of elements following `DISPL` should always be 24 otherwise the `scph`-program will crash.

The simplest way to obtain the parameters `NDISP` and `DISPL` is to generate the supercell with the `phon`-program, i.e by setting `LSUPER = .T.` in the `INPHON`-file and to run the `phon`-program. The output of such a run will not only generate the `SPOSCAR`-file but also a `DISP`-file.

Example 1:

If the initial `POSCAR`-file used to generate the `SPOSCAR`-file was that of a primitive bcc -cell (see `exaples/bcc/`), the following `DISP`-file would be generated:

```
1 0.00000000 0.00100000 0.00100000
```

revealing that we only need one initial displacement, i.e have to move the first atom in the `SPOSCAR`-file and this atom should be moved by the vector

```
0.00000000 0.00100000 0.00100000.
```

The corresponding line 2 and 3 in the `INPUTFLJ`-file would then read:

```
NDISP 1
```

```
DISPL 1 0.0 0.001 0.001  n x.x y.y z.z  n x.x y.y z.z  n x.x y.y z.z  ....
```

where `n x.x y.y z.z....` is used to indicate the arbitrariness of the second to sixth block of the array `DISPL`.

Example 2:

If the initial POSCAR-file used to generate the SPOSCAR-file was that of a primitive hcp-cell (see `examples/hcp/`), the following DISP-file would be generated:

```
1 0.00120000 0.00120000 0.00000000
1 0.00000000 0.00000000 0.00200000
```

revealing that two displacements had to be used in order to obtain the initial phonon spectrum. The corresponding line 2 and 3 in the INPUTFLJ-file would then read:

```
NDISP 2
DISPL 1 0.0012 0.0012 0.0 1 0.0 0.0 0.002 n x.x y.y z.z n x.x y.y z.z....
```

Also do not forget that the supercell files `POSCAR_START1`, `POSCAR_START2`, ... used in the calculation of the initial phonon frequencies should be obtained through modifications of the SPOSCAR-file according to line 3 in the INPUTFLJ-file. For example see the files `examples/bcc/POSCAR_START` and `examples/hcp/POSCAR_START1`, `examples/hcp/POSCAR_START2`.

Line (4-19):

On this lines of the INPUTFLJ-file the different model-potential parameters are specified. Later in Chapter 4 these parameters will be related to existing model-potentials. WARNING, even though you decide to calculate the inter-atomic forces externally, i.e you set the flag `POT` equal to `ABI` the lines 4-19 must still be left in the INPUTFLJ-file otherwise the `scph` program will crash. The safest approach is to use one of the INPUTFLJ-files stored in the `examples/`-directory as a Generic INPUTFLJ-file and modify it according to your needs.

Line (20-21):

The parameter `SAMPLING` at line 20 determines if the atomic displacement amplitudes, $\mathcal{R}_{\mathbf{q}s}$, should be approximated by a discretized version of a gaussian distribution, or if they should be sampled from the true gaussian distribution. If the parameter `SAMPLING` is equal to `GAUSS` then the sampling of the atomic displacement amplitudes, $\mathcal{R}_{\mathbf{q}s}$, will be gaussian with mean square deviation, σ , equal to

$$\sigma^2(\omega_{\mathbf{q},s}, T) = \frac{\langle Q_{\mathbf{q}s}^\dagger Q_{\mathbf{q}s} \rangle}{M} = \frac{\hbar}{M\omega_{\mathbf{q}s}} \left[\frac{1}{2} + n \left(\frac{\hbar\omega_{\mathbf{q}s}}{k_B T} \right) \right]. \quad (3.2)$$

If the parameter `SAMPLING` is not equal to `GAUSS`, then the atomic displacement amplitudes, $\mathcal{R}_{\mathbf{q}s}$, will be sampled from discrete set of amplitudes $\pm\mathcal{R}_1, \pm\mathcal{R}_2, \dots, \pm\mathcal{R}_n$, where the number, n , of amplitudes is given by the input parameter `NGAUSSAMPLE` on line 21. The discrete probability distribution $\mathcal{P}(\pm\mathcal{R}_i)$, $i = 1, \dots, n$ is then constructed in such a way that

$$2 \sum_i^n \mathcal{P}(\mathcal{R}_i) \mathcal{R}_i^2 = \sigma^2(\omega_{\mathbf{q},s}, T) \quad (3.3)$$

is always fulfilled.

Line(22):

This line contains the parameter `MAXITTER` and in it the maximum number of SCAILD iterations is stored. This parameter is used to avoid that a periodic sequences of +/- signs will appear when the Fourier-amplitudes $\mathcal{R}_{\mathbf{q}_s}$ given in Eqn. 2.9 are calculated, i.e the parameter is used to assure that the +/- are chosen in a "truly" random way.

Line(23-25):

The parameter `DSITTER` is the number of iterations up to which the `scph`-code in the calculations of the atomic displacements will replace commensurate phonon frequencies whenever the frequencies are *too close to zero* with:

$$\omega_{\mathbf{q}_s} = \omega_D \frac{|\mathbf{q}|}{|\mathbf{q}_{max}|}. \quad (3.4)$$

Here ω_D is the Debye frequency, \mathbf{q} the wave-vector of the phonon mode and $|\mathbf{q}_{max}|$ is the distance to the Brillouin-zone boundary in the direction of \mathbf{q} . This is a safety mechanism implemented to avoid singular displacements during the first `DSITTER`-iterations of the self consistent cycle. *To close to zero* is defined by the parameter `MAXAMP` (line 25, in units of Å), i.e the above replacement (Eqn. 3.4) only take place whenever $\mathcal{R}_{\mathbf{q}_s} > \text{MAXAMP}$ and the number of iterations is less than `DSITTER` (for a definition of $\mathcal{R}_{\mathbf{q}_s}$ see Eqn. 2.8 - 2.9). However, if the parameter `SUPERSAFE` (line 24) is set to `.TRUE.`, then whenever $\mathcal{R}_{\mathbf{q}_s} > \text{MAXAMP}$ and number of iterations $< \text{DSITTER}$ the mode contribution to the atomic displacements, $\mathcal{R}_{\mathbf{q}_s}$, is set to zero.

3.2.4 The DIPOLEINTERACT-file

If this file is present in the directory where the `scph`-program is executed, the `scph`-program will use the data stored in the `DIPOLEINTERACT`-file to adjust the Longitudinal-optical and Transverse-optical phonon frequencies at the Γ -point for the long-ranged dipole-dipole interaction. The non-analytical dipole-dipole interaction correction to the dynamical matrix in the vicinity of the Γ -point is given by the expression found in Parlinski *et al* [4]

$$\mathcal{D}_{mn}^{\mu\nu}(\mathbf{q})_{corr} = \frac{4\pi e^2}{V \sqrt{M_\mu M_\nu}} \frac{(\bar{\bar{Z}}^\mu \mathbf{q})_m (\bar{\bar{Z}}^\nu \mathbf{q})_n}{\mathbf{q} \bar{\bar{\epsilon}}_\infty \mathbf{q}} e^{-i\mathbf{G}(\mathbf{b}_\mu - \mathbf{b}_\nu)} e^{-\mathbf{q}^2/\rho^2}. \quad (3.5)$$

Here m, n are cartesian coordinate indices, V the volume of the primitive cell, M_μ , $\bar{\bar{Z}}^\mu$ is the mass respectively Born-effective-charge matrix of atom μ . Furthermore, $\bar{\bar{\epsilon}}_\infty$ is the static infinite wavelength dielectric matrix, \mathbf{G} is the reciprocal lattice vector at which \mathbf{q} is centered and ρ is a free damping-parameter which is used to suppress the correction term as $|\mathbf{q}| > 0$. Below a `DIPOLEINTERACT`-file is given as an example

```

0.25      ← ρ
66.523963392000000000000000000000 ← V
34.0 0.00 0.00 ← (ε∞)1n
0.00 34.0 0.00 ← (ε∞)2n
0.00 0.00 34.0 ← (ε∞)3n
6.16 0.00 0.00 ← Z1nPb
0.00 6.16 0.00 ← Z2nPb
0.00 0.00 6.16 ← Z3nPb
-6.16 0.00 0.00 ← Z1nTe
0.00 -6.16 0.00 ← Z2nTe
0.00 0.00 -6.16 ← Z3nTe

```

This file was used in a `scph`-calculation of PbTe, and the data was taken from the work of Waghmare *et al* [5]. The arrows and text to the right of the arrows are not part of the actual file, they are only printed out for the sake of clarity.

The first and second line of the `DIPOLEINTERACT`-file always contain the free-damping parameter, ρ , and the primitive cell volume, V . The following three lines always correspond to the rows of the dielectric tensor, $\bar{\epsilon}_\infty$. The entries following the rows of the dielectric tensor, are the Born-effective-charge matrices. They are entered in blocks of 3 rows, and the number of blocks (matrices) following the entries of the dielectric matrix should be equal to the parameter value of `NTYPES` found in the `INPHON`-file.

3.2.5 The KRAFTER-file

This file just contains the forces extracted from the `OUTCAR`-file by means of the perl script `force_extract`. The forces stored in this file is used by the `scph`-program to calculate a new set of phonon frequencies by means of Eqn. 2.12. The forces in this file are the forces being calculated in STEP 3 in Fig. 2.1.

3.2.6 The KRAFTER.1, . . . , KRAFTER.n-files

These files contain the forces calculated from the n initial single atom displacements (see STEP 1 in Fig. 2.1). From these forces the initial/starting-guess phonons are calculated. Observe once again that in this version of the `scph`-code the maximum number of initial displacements n are 4.

3.2.7 The QPOINTS and the SYMOP -files

The `QPOINTS`-file contains the q-point mesh used by `scph`-code to calculate the phonon density of states. The `QPOINTS`-file can be generated in to different ways:

(1): Set the parameters `QA`, `QB`, `QC` to the desired q-point resolution and let the `QA`-parameter be positive in the `INPHON`-file and use the `phon`-code to generate the `QPOINTS`-file.

(2): Set the parameters `QA`, `QB`, `QC` to the desired q-point resolution and let the `QA`-parameter be negative in the `INPHON`-file. Make sure that one of the files `QPOINTS` or `SYMOP` is not present in the same directory that you are running the `scph`-code. Then when running the the `scph`-code the `QPOINTS`- and the `SYMOP` -file will be generated when the `scph`-code calculates the initial guess, i.e STEP 1 in Fig. 2.1.

The SYMOP -file contains all the symmetry operations used in Eqn. 2.17. To generate the SYMOP -file make sure that one of the files QPOINTS or SYMOP is not present in the same directory that you are going to run the `scph`-code. Then when the `scph`-code is executed the QPOINTS- and the SYMOP -file will be generated during the initial-guess calculations, i.e during STEP 1 in Fig. 2.1.

3.2.8 The MEMORY-file

If a SCAILD calculation is to be initialized, i.e started from scratch by calculating the initial-guess phonons there is no need to provide the `scph`-code with any MEMORY-file. In fact there should not be any MEMORY-file present in the directory where the `scph`-code is executed if one wants to start a SCAILD calculation from scratch. The MEMORY-file is created as an output-file from the `scph`-code after the first initial calculation. The MEMORY-file contains all the data that needs to be saved in between the external force calculations (STEP 2 and 4 in fig 2.1). The MEMORY-file contains all the phonon eigenvectors $\epsilon_{\mathbf{k}s}$, Fourier amplitudes $\mathcal{R}_{\mathbf{k}s}$ used to calculate the atomic displacements $\mathbf{U}_{\mathbf{R}}$ given by Eqn. 2.5. If the output of $\epsilon_{\mathbf{k}s}$ and $\mathcal{R}_{\mathbf{k}s}$ was stored in the MEMORY-file, at let's say iteration number n , this information is used as input by the `scph`-code the $(n + 1)$:th time the `scph`-code is called, to calculate a new set of frequencies from the forces calculated the $(n + 1)$:th iteration, the stored $\epsilon_{\mathbf{k}s}$ and $\mathcal{R}_{\mathbf{k}s}$ through Eqn. 2.12.

If the SCAILD calculation crashes or is interrupted because the external force calculation crashes or is interrupted, at let's say iteration number $(n + 1)$, and you want to restart the calculation from the n :th iteration, then all you need to do is to make sure that the POSCARTEMP1-file corresponding to the MEMORY-file of the n :th iteration is the same file as the POSCAR-file before you restart the external force calculation (`vasp`).

Furthermore, the topmost entry of the MEMORY-file corresponds to the number of times the `scph`-code has been executed after external force calculations, including the time the `scph`-code has been executed after the "starting-guess" calculation.

3.3 The output files

The output files of the `scph`-program are the following:

```
MEMORYFILE
POSCARTEMP1
DISPLACEMENTS
FREQ
DOS
DOSPROJ
CONVERGENCE
QPOINTS
SYMOP
BLANDAD
```

3.3.1 The MEMORY-file

See previous section 3.2.7 for information about this file.

3.3.2 The POSCARTEMP1-file

The POSCARTEMP1-file contains the direct coordinates of the displaced atoms of the supercell, i.e they contain the positions $\mathbf{R} + \mathbf{U}_{\mathbf{R}}$ where $\mathbf{U}_{\mathbf{R}}$ has been calculated by means of Eqn. 2.5. This file is used to calculate the forces on the atoms in STEP 3 of (see Fig. 2.1) the SCAILD cycle.

3.3.3 The DISPLACEMENTS-file

The DISPLACEMENTS-file contains the atomic displacements $\mathbf{U}_{\mathbf{R}}$.

3.3.4 The FREQ-file

In the FREQ-file the calculated phonon-frequency dispersion along the high-symmetry directions specified in the INPHON-file are stored. The frequencies in this file are the "re-normalized" frequencies $\omega_{\mathbf{k}s}(N_I)$ given in Eqn. 2.18.

3.3.5 The DOS-file

In this file the phonon density of states of the last SCAILD iteration is stored. The phonon density of states is calculated from the "re-normalized" frequencies $\omega_{\mathbf{k}s}(N_I)$ given in Eqn. 2.18, by using the q-point mesh stored in the QPOINTS-file. The phonon density of states is given by

$$g(\omega) = \sum_{\mathbf{q},s} \delta(\omega - \omega_{\mathbf{q}s}). \quad (3.6)$$

3.3.6 The DOSPROJ-file

In this file the atom resolved density of states is stored. The atomic resolved density of states is given by

$$g_{\mu}(\omega) = \sum_{\mathbf{q},s} |\epsilon_{\mathbf{k}s}^{\mu}|^2 \delta(\omega - \omega_{\mathbf{q}s}). \quad (3.7)$$

Here $\epsilon_{\mathbf{k}s}^{\mu}$ is the phonon eigen-vector of the atom type μ .

3.3.7 The CONVERGENCE-file

This file contains information about the progress of the SCAILD calculation. Below the typical output of a CONVERGENCE-file is given:

```

ITTER  F [ eV ]      E [ eV ]      DF [eV]  V[eV] Dos_Integral <U(R)^ 2> [a**2]
  1 -0.3980787615  0.25966453  0.00161204  0.000  0.9991328509  0.0809776567040
  2 -0.3966599357  0.25962036  0.00141882  0.000  0.9999996853  0.0747259537950
  3 -0.4071303956  0.25953731  0.01047045  0.000  0.9999996853  0.0877559584183
  4 -0.4123015394  0.25949212  0.00517114  0.000  0.9999993653  0.0789842087423
  5 -0.4155397163  0.25946007  0.00323817  0.000  0.9999996854  0.0894309802517

```

In the first column the iteration number is given. In the second column the phonon free energy calculated with Eqn. 3.1 is given. In the third column the internal energy is given. In the fourth column the difference in lattice dynamical free energy between consecutive iterations is given. In the fifth column the static lattice energy of the system is given if a model potential is being used to calculate the inter-atomic forces. If the inter-atomic forces are being calculated externally this column will only display zeros. In the sixth column the integrated density of states is given, i.e

$$\frac{1}{3nN} \int_0^\infty g(\omega) d\omega. \quad (3.8)$$

Here n is the number of atoms in the primitive cell and $N = QA \times QB \times QC$. This means that the numbers in the sixth column should be close to 1 in order to assure that no imaginary phonon frequencies exists. The reason why the numbers in the sixth column need not be exactly equal to 1 is because smearing has been employed in the calculation of $g(\omega)$.

In the seventh column the mean square deviation of the atoms from their equilibrium positions are given, i.e

$$\langle U(\mathbf{R}) \rangle \equiv \sum_{\mathbf{R} \in SC} \mathbf{U}_{\mathbf{R}}^2, \quad (3.9)$$

where SC denotes the set of atomic positions of the supercell.

WARNING: If the numbers in the seventh column are $\gtrsim 1 \text{ \AA}$ then the atomic displacements are most probably too big. The reason behind this erratic behavior is that one or several of the phonon frequencies calculated at a commensurate q-point are close to zero. To understand the danger of "commensurate phonon frequencies" being too close to zero please inspect Eqn. 2.8. To avoid this problem try with increasing the integer number of the parameter DSITTER or decreasing the value of the parameter MAXAMP.

3.3.8 The QPOINTS and SYMOP-file

See previous section 3.2.6 for information about these files.

3.3.9 The BLANDAD-file

This file keeps track on whether or not the frequencies corresponding to commensurate q-vectors have been updated since they first were calculated by the starting guess iteration. Below we see a typical output from a BLANDAD-file

```
K= 5
0.0000000000  0.2500000000  0.0000000000
  T
  F
  T
```

The first two lines above give information about the q-vector. The following three lines contains information regarding the individual modes of the q-vector (longitudinal, transverse, optical). If all the letters following the first two lines are T it means that the `scph`-program has successfully managed to project out the frequency of all the modes of the q-vector (see Eqn. 2.12). If the letter F appears on any of these lines it means that the frequency of at least one of the modes has not been updated since the initial starting-guess calculation.

Chapter 4

Model-potential calculations

As has been mentioned previously in this manual there is also the possibility to calculate the inter-atomic forces internally, i.e by using the model-potentials implemented within the `scph`-program. In this chapter the analytic formulae of these potentials will be presented and their respective parameters will be coupled to the data in the `INPUTFLJ`-file.

4.1 The MSV-potential

By setting the input parameter `POT` in the `INPUTFLJ`-file to `MSV` the inter-atomic forces will be calculated from the *Morse-spline-van der Waals* analytical pair potential [18], given by

$$V(R) = \epsilon \begin{cases} e^{-2\beta(x-x_1)} - 2e^{-\beta(x-x_1)}, & 0 < x \leq x_1 \\ b_1 + (x - x_1)\{b_2 + (x - x_2)[b_3 + (x - x_1)b_4]\}, & x_1 < x < x_2 \\ -C_6x^{-6} - C_8x^{-8} - C_{10}x^{-10}, & x_2 < x \end{cases} \quad (4.1)$$

where $x \equiv R/R_m$, and R_m being the interatomic separation at the minimum of the potential. In the following the above parameters will be coupled to their respective input in the `INPUTFLJ`-file. In the list below the model-potential parameters will be presented to the left of the equality signs, and their corresponding input parameters will be presented to the right of the equality sign. (The same convention will be used for all the following model-potentials)

$R_m = \text{Rm}$

$\beta = \text{BETA}$

$x_1 = \text{X1}$

$x_2 = \text{X2}$

$b_1 = \text{B1}$

$b_2 = \text{B2}$

$b_3 = \text{B3}$

$b_4 = \text{B4}$

$C_6 = \text{C6}$

$C_8 = \text{C8}$

$C_{10} = \text{C10}$

$\epsilon = \text{EPS}$

4.2 The ZRP-potential

By setting the input parameter POT in the INPUTFLJ-file to ZRP the inter-atomic forces will be calculated from the N-body analytical pair potential given by Willaime et al [19], given by

$$V(R) = A \sum_{R < R_c} e^{\left[-p \left(\frac{R}{R_0} - 1\right)\right]} - \left\{ \xi^2 \sum_{R < R_c} e^{\left[-2q \left(\frac{R}{R_0} - 1\right)\right]} \right\}^{1/2} \quad (4.2)$$

In the following the above parameters will be coupled to their respective input in the INPUTFLJ-file.

$A = \text{B1}$
 $\xi = \text{B2}$
 $p = \text{B3}$
 $q = \text{B4}$
 $R_0 = \text{X1}$
 $R_c = \text{X2}$

4.3 The EAM-potential

By setting the input parameter POT in the INPUTFLJ-file to EAM the inter-atomic forces will be calculated from the embedded atom method (EAM) type of potential [20]. For this potential type the total potential energy is given by:

$$E = \sum_i \left[\sum_j \phi(R_{ij}) + \sqrt{\sum_j h^2(R_{ij})} \right], \quad (4.3)$$

where

$$\phi(R) = A e^{-p \left(\frac{R}{R_0} - 1\right)} e^p \left[-\left(\frac{R}{R_c}\right)^{n_c} + \left(\frac{R_0}{R_c}\right)^{n_c} \right], \quad (4.4)$$

$$h(R) = B e^{-q \left(\frac{R}{R_0} - 1\right)} e^q \left[-\left(\frac{R}{R_c}\right)^{n_c} + \left(\frac{R_0}{R_c}\right)^{n_c} \right]. \quad (4.5)$$

Here the index j is that of an atom at distance R_{ij} from the atom i . In the following the above parameters will be coupled to their respective input in the INPUTFLJ-file.

$A = \text{B1}$
 $B = \text{B2}$
 $p = \text{B3}$
 $q = \text{B4}$
 $R_0 = \text{X1}$
 $R_c = \text{X2}$
 $n_c = \text{Rm}$

4.4 The PAS-potential

By setting the input parameter POT in the INPUTFLJ-file to PAS the inter-atomic forces will be calculated from the embedded atom method (EAM) type of potential [21]. For this potential type the total potential energy is given by:

$$E = \sum_i \left[\frac{1}{2} \sum_{j(\neq i)} V(R_{ij}) + F(\rho_i) \right]. \quad (4.6)$$

Here the pair potential part is given by

$$V(r) = \sum_{k=1}^7 A_k (R_k - r)^3 \Theta(R_k - r), \quad (4.7)$$

where $\Theta(x > 0) = 1$, $\Theta(x < 0) = 0$. The density is given by

$$\rho_i = \sum_{j(\neq i)} \frac{R_{ij}}{a} e^{-\frac{\beta R_{ij}}{a}}, \quad (4.8)$$

and the embedding function correct to second order in ρ is given by

$$F(\rho) = F_0 + \frac{F_0''}{2} \rho^2 \quad (4.9)$$

In the following the above parameters will be coupled to their respective input in the INPUTFLJ-file.

$R_k = \text{RK}$

$A_k = \text{AK}$

$\beta = \text{BETA}$

$a = \text{Rm}$

$F_0'' = \text{X1}$

$F_0 = 0$ (you don't have to set this parameter)

4.5 The PIN-potential

By setting the input parameter POT in the INPUTFLJ-file to PIN the inter-atomic forces will be calculated from the embedded atom method (EAM) type of potential [22]. For this potential type the total potential energy is given by:

$$E = \sum_i \left[\frac{1}{2} \sum_j V(R_{ij}) + \sqrt{\rho_i} \right]. \quad (4.10)$$

Here the pair potential part is given by

$$V(r) = \sum_{k=1}^7 A_k (R_k - r)^3 \Theta(R_k - r), \quad (4.11)$$

and the density is given by

$$\rho_i = \sum_j \sum_k^2 a_k (r_k - R_{ij})^3 \Theta(r_k - R_{ij}) \quad (4.12)$$

In the following the above parameters will be coupled to their respective input in the INPUTFLJ-file.

R_k = RK (first six numbers from the left in the INPUTFLJ-file)

A_k = AK (first six numbers from the left in the INPUTFLJ-file)

a_1 = B1

a_2 = B2

r_1 = B3

r_2 = B4

4.6 The PAA-potential

By setting the input parameter POT in the INPUTFLJ-file to PAA the inter-atomic forces will be calculated from the embedded atom method (EAM) type of potential [23]. For this potential type the total potential energy is given by:

$$E = \sum_i \left[\frac{1}{2} \sum_{j(\neq i)} V(R_{ij}) + F(\rho_i) \right]. \quad (4.13)$$

Here the pair potential part is given by

$$V(r) = \sum_{k=1}^7 A_k (R_k - r)^3 \Theta(R_k - r), \quad (4.14)$$

and the density is given by

$$\rho_i = \sum_j \phi_0 f(R_{ij}), \quad (4.15)$$

where

$$f(r) = \begin{cases} \frac{e^{-5r}}{r}, & r \leq R_m \\ (r - R_c)^3 (a_1 r^2 + a_2 r + a_3), & R_m < r \leq R_c \\ 0, & R_c \leq r \end{cases} \quad (4.16)$$

Furthermore the embedding function for the EAP of Zr is expressed as a 8:th order polynomial

$$F(\rho) = \sum_{k=0}^8 B_k \rho^k. \quad (4.17)$$

In the following the above parameters will be coupled to their respective input in the INPUTFLJ-file.

$R_k = \text{RK}$
 $A_k = \text{AK}$
 $R_m = \text{X1}$
 $R_c = \text{X2}$
 a (lattice constant of reference structure in Ref. [23]) (= Rm)
 $\phi_0 = \text{EPS}$
 $a_1 = \text{AZ}$ (first element, counting from the left in the INPUTFLJ-file, i.e. $\text{AZ}[1]$)
 $a_2 = \text{AZ}[2]$
 $a_3 = \text{AZ}[3]$
 $B_8 = \text{B1}$
 $B_7 = \text{B2}$
 $B_6 = \text{B3}$
 $B_5 = \text{B4}$
 $B_4 = \text{C6}$
 $B_3 = \text{C8}$
 $B_2 = \text{C10}$
 $B_1 = \text{AZ}[4]$
 $B_0 = \text{BETA}$

Bibliography

- [1] P. Souvatzis, O. Eriksson, M. I. Katsnelson and S. P. Rudin, Phys. Rev. Lett **100**, 095901 (2008).
- [2] G. Kresse & J. Furthmuller, Phys. Rev. B **54**, 11169 (1996).
- [3] The phon program written by Dario Alfe can be obtained from <http://chianti.geol.ucl.ac.uk/~dario/>
- [4] K. Parlinski, Z. Q. Li, and Y. Kawazoe, Phys. Rev. Lett. **81**, 3298 (1998)
- [5] U. V. Waghmare, N. A. Spaldin, H. C. Kandpal, and Ram Seshadri, Phys. Rev. B **67**, 125111 (2003)
- [6] A. Heiming, W. Petry, J. Tramenau, M. Alba, C. Herzig, H. R. Schober, and G. Vogl. Phys. Rev. B **43**, 10948 (1991).
- [7] W. Petry, A. Heiming, J. Tramenau, M. Alba, C. Herzig, H. R. Schober, and G. Vogl. Phys. Rev. B **43**, 10933 (1991).
- [8] J. Tramenau, A. Heiming, W. Petry, M. Alba, C. Herzig, W. Miekeley, and H. R. Schober, Phys. Rev. B **43**, 10963 (1991).
- [9] W. Petry, J. Tramenau and C. Herzig, Phys. Rev. B **48**, 881 (1993).
- [10] M. I. Katsnelson, A. V. Trefilov, and K. Yu. Khromov, JETP Lett. **69**, 688 (1999).
- [11] M. I. Katsnelson, A. V. Trefilov, M. N. Khlopkin, and K. Yu. Khromov, Phil. Mag. B **81**, 1893 (2001).
- [12] M. van Schilfgaarde, T. Kotani, and S. Faleev, Phys. Rev. Lett. **96**, 226402 (2006)
- [13] P. Souvatzis, A. Delin, and O. Eriksson, Phys. Rev. B **73**, 054110-054116 (2006).
- [14] M. I. Katsnelson, A. F. Maksyutov, and A. V. Trefilov, Phys. Lett. A **295**, 50 (2002).
- [15] C. Stassis, D. Arch, B. N. Harmon and N. Wakabayashi, Phys. Rev. B **19**, 181-188 (1979).
- [16] C. Stassis, J. Zarestky, D. Arch, O. D. McMasters and B. N. Harmon, Phys. Rev. B **18**, 2632 (1978).

- [17] C. Stassis, D. Arch, O. D. McMasters and B. N. Harmon, Phys. Rev. B **24**, 730-740 (1981).
- [18] R. N. Kersarwani, Y. P. Varshni, Phys. Rev. A **23**, 991 (1981).
- [19] F. Willaime and C. Massobrio, Phys. Rev. Lett. **63**, 2244 (1989).
- [20] K. Masuda-Jindo, S. R. Nishitani and Vu VanHung, Phys. Rev. B **70**, 184122 (2004).
- [21] R. Pasianot and E. J. Savino, Phys. Rev. B **45**, 12704 (1992).
- [22] U. Pinsook and G. J. Ackland, Phys. Rev. B **58**, 11252 (1998).
- [23] R.C Pasianot and A.M. Monti, J. Nucl. Mat. **264**, 198-205 (1999).